

REGULAR MINIMAL NETS ON SURFACES OF CONSTANT NEGATIVE CURVATURE

ALINA VDOVINA

Khmel'nitskiy Thechnical University

e-mail: alina@alpha.podol.khmel'nitskiy.ua

E.N. SELIVANOVA*

Department of Geometry, Nizhny Novgorod State Pedagogical University

603000 Russia, Nizhny Novgorod, ul. Ulyanova 1

e-mail: lena@moebius.mathematik.uni-tuebingen.de

Abstract. The problem of classification of closed local minimal nets on surfaces of constant negative curvature has been formulated in [3], [4] in the context of the famous Plateau problem in the one-dimensional case. In [6] an asymptotic for $\log \sharp(W^r(g))$ as $g \rightarrow +\infty$ where g is genus and $W^r(g)$ is the set of regular single-face closed local minimal nets on surfaces of curvature -1 has been obtained. It has been shown that $\sharp(W^r(g))$ is equal to the number of classes of topological equivalence of all single-face closed local minimal nets on surfaces of curvature -1 . In this paper we prove an asymptotic for $\sharp(W^r(g))$ as $g \rightarrow +\infty$ and construct some examples of f -face, $f > 1$ nets.

1 Introduction

Let (M^2, ds^2) be a Riemann surface of constant curvature. *Closed local minimal net* Γ on M^2 is an embedded cubic graph consisting of segments of geodesics where the angles at any vertex are equal to $2\pi/3$. A local minimal net is *regular* if all segments of geodesics are equal.

Two local minimal nets are *topologically equivalent* if there is a homeomorphism of surfaces preserving the orientation and transferring nets one into another.

Two local minimal nets are *isomorphic* if there is an isometry of surfaces transferring these nets one into another.

Let Γ be a net on M^2 . If the difference $M^2 \setminus \Gamma$ is homeomorphic to one disk then the net Γ is *single-face*. If the difference $M_g^2 \setminus \Gamma$ is homeomorphic to f disks then net Γ is *f-face*.

The problem of classification of closed local minimal nets on surfaces of constant negative curvature has been formulated in [3] in the context of the famous Plateau problem in the one-dimensional case.

*Supported by DAAD.

Full description of such nets in the case of curvature 1 one can find in [2]. There are many interesting applications of this result, see [1], [3], [4].

We consider the case of orientable surfaces of curvature -1 . In [6] an asymptotic for $\log \sharp(W^r(g))$ as $g \rightarrow +\infty$ where g is genus and $W^r(g)$ is the set of regular single-face closed local minimal nets on surfaces of curvature -1 has been obtained. It has been shown that $\sharp(W^r(g))$ is equal to the number of classes of topological equivalence of all single-face closed local minimal nets on surfaces of curvature -1 . In this paper we prove an asymptotic for $\sharp(W^r(g))$ as $g \rightarrow +\infty$ and show that a description of all single-face closed minimal nets on surfaces of curvature -1 can be obtained due to a description of the set of regular single-face closed minimal nets in terms of polygons on the hyperbolic plane. Then we construct some examples of f -face nets, $f > 1$.

2 Single-face nets

In order to describe the set of the single-face regular local minimal nets on M_g^2 will use the following algebraic construction.

Let us call a word U of length $12g - 6$ a Wicks form of genus g if it satisfies the following conditions

1. each letter has been used twice as x and x^{-1} ,
2. there is no pair of distinct, noninverse letters x, y which appear in U only in subwords $(xy)^{\pm 1}$.

If letters x, y appear in U only in subwords $(xy)^{\pm 1}$, we can replace $(xy)^{\pm 1}$ by $z^{\pm 1}$. This operation we will call *reduction*.

Two words U_1 and U_2 are *equivalent*, if U_1 can be obtained from U_2 by permutation of letters, by substitution of some of them for inverse ones and by cyclic permutation.

It turns out that there is an *inductive method* to construct all words $U(g+1)$ from $U(g)$. We can describe the *transformations* by which all words $U(g+1)$ can be obtained from words $U(g)$.

Let us rewrite U changing a into a_1a_2 and put the fragment

$$id^{-1}ef^{-1}dhe^{-1}fh^{-1}i^{-1}$$

between a_1 and a_2 or between a_2^{-1} and a_1^{-1} . This transformation is called α -transformation.

Another transformation is to rewrite U changing a into a_1a_2 and b into b_1b_2 and put the fragment

$$d^{-1}ef^{-1}d$$

between a_1 and a_2 or a_2^{-1} and a_1^{-1} and put the fragment

$$he^{-1}fh^{-1}$$

between b_1 and b_2 or b_2^{-1} and b_1^{-1} . This transformation is called by β -transformation.

The third transformation is the following. We rewrite U changing a into a_1a_2 , b into b_1b_2 and c into c_1c_2 i.e.

$$U' = U_1a_i^\epsilon a_j^\epsilon U_2b_k^\delta b_l^\delta U_3c_m^\psi c_p^\psi U_4$$

(an expression of type $a_i^\epsilon a_j^\epsilon$ means the a_1a_2 or $a_2^{-1}a_1^{-1}$, where $\epsilon = 1, i = 1, j = 2$ or $\epsilon = -1, i = 2, j = 1$) We'll say that the word

$$V = U_1a_i^\epsilon d^{-1}eb_l^\delta U_3c_m^\psi f^{-1}da_j^\epsilon U_2b_k^\delta e^{-1}fc_p^\psi U_4$$

is obtained from U by γ -transformation.

Lemma, [5]. Word $U(g)$ corresponds to an embedded cubic graph with $4g - 2$ vertices on a M_g^2 if and only if this word can be obtained from

$$abca^{-1}b^{-1}c^{-1}$$

by a sequence of $(g - 1)$ α , β or γ -transformations.

Theorem 1.1 For $\sharp W^r(g)$ the following asymptotic

$$\sharp W^r(g) = \frac{(6g - 4)!}{(3g - 2)!g!12^g}(1 + o(1))$$

holds.

Proof. As mentioned above, $\sharp W^r(g)$ equals the number of classes of topological equivalence of the single-face nets on M_g . First we shall show that the problem of topological classification of single-face local minimal nets on $M_g^2, g > 1$ is equivalent to the problem of topological classification of embeddings of cubic graphs to a surface M_g^2 .

Let us consider a single-face embedded cubic graph P in a surface M_g^2 , label the edges of P by different letters $\{a, b, \dots\}$ of an alphabet A . Let us choose also an orientation of any edge of P . The difference $M_g^2 \setminus P$ consists of a disk with a word U on the boundary of this disk. The number of the different letters (or the number of edges of the graph) is $6g - 3$. In the word U each letter has been used twice. Taking into account the orientation of the edges we'll write x and x^{-1} if the orientations of the corresponding segments are different. There is no pair of distinct, noninverse letters x, y which appear in U only in subwords $(xy)^{\pm 1}$. Our classification theorem is based on the description of words U from Lemma before the theorem.

There is a realization of local minimal nets for any equivalence class of cubic graphs on the M_g^2 as a nontrivial subset of this class. (One may consider a regular polygon on the hyperbolic plane labeled according the corresponding word U).

Thus, we have now to describe the set of equivalence classes of graphs. This means to describe all possible words U up to equivalence, see above.

Let us consider an oriented connected cubic graph Γ with $2k$ vertices and $3k$ edges labeled by different letters of an alphabet A with specified cyclic orderings of the edges at any vertex. At any vertex v of graph we will write a cyclic ordered

triples of letters so that a letter x has exponent 1 if the edge x enters in v and -1 if x leaves v . An *ordered graph* is a graph with specified cyclic orderings (the triples) at the edges at each vertex. We will call the set of elements $x, x^{-1}, x \in A$ by $2Q$. The vertices of a graph may be considered as sets into which $2Q$ is partitioned. Let us consider a permutation P on $2Q$ such that the cyclic order at any vertex determines a cycle in the permutation P and an inverse ξ transferring x to x^{-1} for any x from $2Q$. If permutation $P\xi$ has f cycles then permutation P determines an embedding of graph Γ in the sphere with g handles where there are f faces, $2k$ vertices and $3k$ edges, satisfying the equality $k = 2g + f - 2$, [7], i.e. there is an f -face embedding of genus g . (Note that for $f = 1$ permutation $P\xi$ is the considered above Wicks form U). Two ordered graphs (Q, P) and (Q', P') are *isomorph* if there is a one-to-one mapping Φ such that

$$\Phi[P(\beta)] = P'[\Phi(\beta)], \Phi(\beta^{-1}) = \Phi(\beta)^{-1}$$

for any β from $2Q$. Thus the *automorphism* group of P (denoted by $aut(P)$) is the set of permutations on $2Q$ which commute with P and ξ .

A connected ordered graph is rooted if one element β of $2Q$ is distinguished. A *root isomorphism* from one ordered graph (E, P, β) onto another one (E', P', β') is an isomorphism Φ from (E, P) onto (E', P') such that $\Phi(\beta) = \beta'$. Thus a *root automorphism* on (E, P, β) is a permutation on $2E$ which commutes with P and ξ and preserves β . (Corresponding Wicks form we will call a rooted Wicks form.) In [7] has been shown that *the number of essentially different ways to root the ordered graph (E, P)* that is, the number of root-isomorphism classes of rooted ordered graphs which are isomorphic as ordered graphs to (E, P) is $6k/|aut(P)|$. Then

$$1 \leq t \leq 6k. \quad (1)$$

Let $R(1, g)$ be the number of rooted ordered graphs on $4g - 2$ vertices, $6g - 3$ edges such permutation $P\xi$ consists of one cycle.

The following formula has been obtained in [7]:

$$R(g) = \frac{(12g - 6)(6g - 4)!}{(3g - 2)!g!12^g}. \quad (2)$$

Let $N(1, g)$ be the number of rooted ordered graphs on $4g - 2$ vertices, $6g - 3$ edges such permutation $P\xi$ consists of one cycle. Then $R(g)/t \leq N(1, g) \leq R(g)$. There is a one-to-one correspondence between single-face genus g embeddings and ordered graphs where permutation $P\xi$ consists of one $12g - 6$. Then the number of genus g embeddings equals to $N(1, g)$. Taking into account (1) and (2) we obtain

$$\begin{aligned} \frac{(6g - 4)!}{(3g - 2)!g!12^g} &\leq N(1, g) \leq \\ &\leq \frac{(12g - 6)(6g - 4)!}{(3g - 2)!g!12^g}. \end{aligned} \quad (3)$$

Thus

$$N(1, g) \geq \frac{R(g)}{12g - 6}.$$

Let us prove $N(1, g) \leq \frac{R(g)}{12g - 6}(1 + o(1))$.

Denote $N_1(g)$ the number of the Wicks forms with trivial automorphisms and $N_k(g)$ the number of the Wicks forms with automorphisms of order k . It is obvious, that

$$N_1(g) \leq R(g)/12g - 6.$$

So, we have to prove, that

$$\sum_{k=2}^{12g-6} N_k(g) = o(R(g)/12g - 6). \quad (1)$$

Let x and y be some letters of U where $U = U_1xU_2y$. We define the distance between x and y (it is noted as $d(x, y)$) for letters x, y of U to be $\min(|U_1|, |U_2|)$. We will estimate the number of boundary walks of single-face embeddings of cubic graphs on $4g - 2$ vertices. It is obvious, that a boundary walk of a single-face embedding of a cubic connected graph on $4g - 2$ vertices can be written as a word U with length $12g - 6$. The word U will be a genus g Wicks form. Let us note, that if edges a, b, c are leaving a vertex v of G , then there are subwords $a^{-1}b, b^{-1}c$ and $c^{-1}a$ in U .

Lemma 1. One may estimate the number of the Wicks forms with automorphisms of the second order by the following expression

$$N_2(g) \leq (g + 1) \prod_{m \in N, 12g-6-7m > 0} (12g - 6 - 7m).$$

Proof. We will consider a $12g - 6$ -gon and try to construct a Wicks form U with an automorphism ξ of the second order. It is known, (see, for example [5]), that every multiple edge (b, c) gives subwords of $abca^{-1}$ and $db^{-1}c^{-1}d^{-1}$ kind in U or subword of $eabca^{-1}db^{-1}c^{-1}d^{-1}e^{-1}$ kind. In the beginning of the proof we consider the case, when U has l multiple edges such that $db^{-1}c^{-1}d^{-1} = \xi(abca^{-1})$. We will say, that the multiple edge (b, c) is *diagonal multiple edge*.

On the first step we fix words $a_1b_1c_1a_1^{-1}$ and $d_1b_1^{-1}c_1^{-1}d_1^{-1}$ corresponding to a multiple edge (b_1, c_1) , $d_1b_1^{-1}c_1^{-1}d_1^{-1} = \xi(a_1b_1c_1a_1^{-1})$. So, 8 letters are fixed before the second step and there are no more than $12g - 6 - 8$ possibilities for the words corresponding to the second multiple edge (a_2, b_2) . There are 16 letters are fixed before the third step and there are no more than $12g - 6 - 16$ possibilities for the words corresponding to the third multiple edge. The procedure we will call α - procedure and $8k$ letters have been already fixed after k -step of the α - procedure. The α - procedure has l steps and there are $12g - 6 - 8l$ "free" places after it. Let's describe a β -procedure. Let's note, that if letter x is fixed, then x^{-1} , $\xi(x)$ and $\xi(x^{-1})$ are fixed too. Let z be a letter, which have been already fixed, but there is a free place near z . We continue l -step of α - procedure by fixing elements x_1

and y_1 such that $z^{-1}x_1$ and $y_1^{-1}z$ are subwords of U and x_1 and y_1 have not been fixed by α - procedure. So, there are no more, than $12g - 6 - 8l - 2$ possibilities for the subword $x_1^{-1}y_1$ in U . The places for $\xi(x_1), \xi(y_1), \xi(x_1^{-1}), \xi(y_1^{-1})$ are determined, because U has an automorphism of the second order. Let w be a letter, which has already fixed, but there are free places after w and before w^{-1} . We will put x_2 after w and y_2 before w^{-1} . We define k -step of the β -procedure so that elements $x_k, y_k, \xi(x_{k-1}), \xi(y_{k-1}), x_{k-1}^{-1}y_{k-1}^{-1}, \xi(x_{k-1}^{-1}), \xi(y_{k-1}^{-1})$ are fixed on the k -step. Let A_k be the set of all letters of U which have been already fixed after the k -step. So, $|A_k| = |A_{k-1}| + |N(k)|$, where $N(k)$ is a number of letters, which can be fixed on the k -step. By definition of k -step, the set $N(k)$ consists of the elements $x_k, y_k, \xi(x_{k-1}), \xi(y_{k-1}), x_{k-1}^{-1}y_{k-1}^{-1}, \xi(x_{k-1}^{-1}), \xi(y_{k-1}^{-1})$. According our construction these elements has been not fixed before k -step. It easy to see, that $N(k) \geq 7$, if x_{k-1} and y_{k-1} are not multiple edges with $x_{k-1} = \xi(x_{k-1}), y_{k-1} = \xi(y_{k-1})$.

We will continue the β -procedure while it is possible. So, the general procedure of constructing U consists of two inductive procedures(α and β). We will say, that the procedure of constructing of a cubic with l diagonal multiple edges is *successful*, if we obtain a Wicks form with exactly l diagonal multiple edges after it. Since we have no more than $12g - 6 - 7k$ possibilities of choice in the k step of the general procedure and no less than seven letters are fixed in the each step, then

$$N^l \leq (12g - 6 - 8) \prod_{m \geq 2, 12g-6-7m > 0} (12g - 6 - 7m), l = 1, \dots, g,$$

where N^l is the number of cubic words with an automorphism of the second order with l diagonal multiple edges (it is easy to see, that a cubic word cannot have more than g multiple edges), $l = 0, \dots, g$.

If U has no multiple edges, we can start β -procedure with a letter c such that $d(c, c^{-1}) = 6g - 4$. Since $\text{Aut}(U) = 2$ the letter c there exists. In this case in the first step we fix only six letters(in the k -step, where $k \geq 2$ we fix no less than seven letters by definition of the β -procedure),

$$N^0 \leq (12g - 6 - 6) \prod_{m \geq 2, 12g-6-7m > 0} (12g - 6 - 7m).$$

So,

$$N_2(g) = N^0 + N^1 + \dots + N^g \leq (g + 1) \prod_{m \in N, 12g-6-7m > 0} (12g - 6 - 7m)$$

The Lemma 1 is proved.

Lemma 2. One may estimate the number of the Wicks forms with automorphisms of order 3 by the following expression

$$N_3(g) \leq 2g^2 \prod_{m \in N, 12g-6-7m > 0} (12g - 6 - 7m).$$

Proof. Let U be a Wicks form with automorphism ξ of the third order. Let v be a vertex and edges a, b, c are leaving it. We will say, that v is automorphic, if $\xi(a^{-1}b) = b^{-1}c$ and $\xi(b^{-1}c) = c^{-1}a$ or $\xi(a^{-1}b) = c^{-1}a$ and $\xi(c^{-1}a) = b^{-1}c$.

Let v_1 be an automorphic vertex, edges a_1, b_1, c_1 are leaving it and the edge b_1 enters a vertex w_1 . It is obvious, that w_1 cannot be automorphic. Edges b_1, b_2, b_3 enter the vertex w_1 and b_2 leaves a vertex v_2 . The vertex v_2 can be automorphic. If v_2 is automorphic, we will say, that v_2 is *neighbor* of v_1 . So, two automorphic vertices v and w are *neighboring*, if the shortest path between v and w consists of two edges.

Vertices, which are ends of an edge x we will denote by x_- and x_+ , where x leaves x_- and enters x_+ . Let us prove, if $|U| \geq 30(g \geq 3)$, then every automorphic vertex can have only one neighbor. Let edges a, b, c be leaving a vertex v , edges d, f, e be leaving a vertex w and v and w are neighboring automorphic vertices, then a_+, b_+, c_+ and d_+, f_+, e_+ are equal sets of vertices of Γ , that is if $b_+ = e_+$, then $a_+ = d_+, c_+ = f_+$ or $a_+ = f_+, c_+ = e_+$. Without loss of generality we can assume, that $a_+ = d_+, b_+ = e_+$ and $c_+ = f_+$. Since all vertices of the graph Γ are degree three, there are some edges g, h, i such that $a_+ = d_+ = g_+, b_+ = e_+ = h_+$ and $c_+ = f_+ = i_+$. If g_+ is automorphic, then $g_+ = h_+ = i_+$ and Γ has only three vertices. So, if $|U| \geq 30(g \geq 3)$, then every automorphic vertex can have only one neighbor.

Let $U = U_1 U_2 U_3$ be a Wicks form with automorphism ϕ of the third order with t single automorphic vertices (without neighbors) and s pairs of automorphic vertices, such that $|U_1| = |U_2| = |U_3|$ and $U_2 = \phi(U_1), U_3 = \phi^2(U_1)$.

Now we describe a transformation of ψ -type. Let $U_1 = x_1 x_2 \dots x_k$. First, we delete all letters from all automorphic vertices. Then, if both x_i and x_i^{-1} belong to U_1 , then we leave x_i and x_i^{-1} without changing. If x_i belongs to U_1 , but x_i^{-1} belongs to U_2 or U_3 , we replace the image of x_i^{-1} in U_1 by x_i^{-1} . After all reductions and cancellations we will obtain a word W , which is a Wicks form of genus p , where $p \leq (g + 2)/3$. We will say, that W is obtained from U by a transformation of ψ -type. On the other hand, there are no more, than

$$3^{12p} \frac{(12p - 6)(12p - 2) \dots (12p - 6 + 4s)(12p - 2 + 4s) \dots (12p - 6 + 4s + 4t)}{s!t!}$$

possibilities to obtain a genus g Wicks form of the third with t single automorphic vertices and s pairs of automorphic vertices from every rooted Wicks form of genus p . The number of nonequivalent rooted cubic words is

$$\frac{(12p - 6)(6p - 4)!}{(3p - 2)!p!12^p}.$$

So, the number of genus g Wicks forms of the third with t single automorphic vertices and s pairs of automorphic vertices, is no more, than

$$S(p) = 3^{12p} (12p - 6)(12p - 2) \dots (12p - 6 + 4s)(12p - 2 + 4s) \dots \\ \frac{(12p - 6 + 4s + 4t)(12p - 6)(6p - 4)!}{s!t!(3p - 2)!p!12^p}.$$

It is easy to see, that

$$S(p) \leq 2g^2 \prod_{m \in N, 12g-6-7m > 0} (12g - 6 - 7m).$$

Lemma 3. One may estimate the number of the Wicks forms with automorphisms of order 5 by the following expression

$$N_5(g) = o(R(g)/12g - 6).$$

Proof. It is easy to see, that a Wicks form with automorphism of order five cannot have any "diagonal" edges and automorphic vertices. So, every vertex v_1 has four different images.

If edges a_1, b_1, c_1 are leaving a vertex v_1 , then there are subwords $a_1^{-1}b_1, b_1^{-1}c_1$ and $c_1^{-1}a_1$ in the Wicks form. Now we will describe an inductive procedure of constructing a genus g Wicks with automorphism of order five. At the first step we fix subwords $a_1^{-1}b_1, b_1^{-1}c_1$ and $c_1^{-1}a_1$ and their images. There are no more than $12g - 6$ possibilities for fixing of subword $a_1^{-1}b_1$, there are no more than $12g - 6 - 2$ possibilities for fixing of subword $b_1^{-1}c_1$ and there are no more than $12g - 6 - 4$ possibilities for fixing of subword $c_1^{-1}a_1$. The places for images of $a_1^{-1}b_1, b_1^{-1}c_1$ and $c_1^{-1}a_1$ are determined and 24 different letters are fixed on the first step by no more than $(12g - 6)(12g - 6 - 2)(12g - 6 - 4)$ possibilities. So, there are no more than $(12g - 6 - 24)$ possibilities for $a_2^{-1}b_2$, no more than $(12g - 6 - 26)$ for $b_2^{-1}c_2$ and no more than $(12g - 6 - 28)$ for $c_2^{-1}b_2$. There are no more than $(12g - 6 - 24)(12g - 6 - 2 - 24)(12g - 6 - 4 - 24)$ variants for fixing 24 letters on the second step. So, on the k -step we fix $a_k^{-1}b_k, b_k^{-1}c_k$ and $c_k^{-1}a_k$ and their images. There are no more than $(12g - 6 - 24(k - 1))(12g - 8 - 24(k - 1))(12g - 10 - 24(k - 1))$ possibilities for this. We have, that a genus g Wicks form can be constructed by no more than

$$\prod_{m \in N, 12g-6-24(m-1) > 0} (12g - 6 - 24(k - 1))(12g - 8 - 24(k - 1))(12g - 10 - 24(k - 1))$$

ways. Lemma 3 follows now after some technical calculations.

Lemma 4. One may estimate the number of the Wicks forms with automorphisms of order k , $k \geq 7$ by the following expression

$$N_k(g) \leq \left\lceil \frac{12g - 6}{7} \right\rceil !.$$

Proof. The proof is obvious.

From the above Lemmas we obtain (1). □

Let us consider an element of the set of topological equivalence of the single-face nets, i. e. a word $U(g) \in W^r(g)$, $U(g) = a...b...c...$, $\{a, b, c, \dots\} = A$ where A is an alphabet. A corresponding local minimal net consists of arcs of geodesics $e_a, e_b, \dots$

Let l_x be length of segment e_x for $x \in A$. Then for any word $U(g)$ there is a system of numbers $L_U = l_a, \dots, l_b, \dots, l_c, \dots$ where $l_x = l_{x^{-1}}$ and the subscripts put together the word $U(g)$.

Thus, the system of numbers defined above is a description of a local minimal net on a surface. On the other hand there are different systems for any net. So, we need to define now an equivalence of these systems. Two systems of numbers $L'_{U'}$ and L_U are equivalent if in some alphabet $U = U'$ and the corresponding components l'_x and l_x are equal for any letter $x \in U$.

What kind of numbers in $\mathbf{R}_+^{12g-6}$ correspond to a system of numbers L_U ? In order to answer on this question we have to describe the set of $12g - 6$ -gons on the hypobolic plane with the following conditions: all angles around vertexes are equal to $2\pi/3$, sides of this polygon can be consequently labeled by letters of the word U so that we can read this word after that and for length l_x of side x holds

$$l_x = l_{x^{-1}}, x \in A. \quad (1)$$

So, there are $6g - 3$ equalities of the type (1) because there are $6g - 3$ different letters in U . Let us now consider three different vertexes A_1, A_2, A_3 of such a $12g - 6$ -gon. We have then three different nonintersect subwords X_1, X_2, X_3 of U . There are analitic functions ν, μ so that

$$\begin{aligned} \hat{A}_1 &= \nu(l_{X_1}, l_{X_2^{-1}}), \\ \hat{A}_2 &= \nu(l_{X_2}, l_{X_3^{-1}}), \\ \hat{A}_3 &= \nu(l_{X_3}, l_{X_1^{-1}}), \\ |A_2 A_3| &= \mu(X_3), \\ |A_1 A_2| &= \mu(X_2), \\ |A_1 A_3| &= \mu(X_1). \end{aligned}$$

It is well known that there is an analitic function ξ such that

$$\begin{aligned} |A_1 A_2| &= \xi(\hat{A}_3, \hat{A}_1, \hat{A}_2), \\ |A_2 A_3| &= \xi(\hat{A}_1, \hat{A}_2, \hat{A}_3), \\ |A_3 A_1| &= \xi(\hat{A}_2, \hat{A}_3, \hat{A}_1). \end{aligned} \quad (2)$$

The equalities (1) and (2) give us a description of all such polygons. The number of conditions is equal to $6g - 3 + 3 = 6g$.

Thus, the set of local minimal nets W_U corresponding to a cubic word U by genus g can be realized as a subset of $\mathbf{R}_+^{12g-6}$ given by equalities (1) and (2).

We obtain also the following theorem.

Theorem 1.2. The set of all closed single-face local minimal nets $W(g)$ can be realized as a union of subsets W_U in $\mathbf{R}_+^{12g-6}$ given by the conditions (1), (2) where U is an element of $W^r(g)$.

Commentar. It is interesting to note that the number of free parameters in the set W_U is equal to $6g - 6$, i. e. the dimension of the Teichmüller space.

3 Exotic nets

Here we suppose some examples of words corresponding to the regular f -face regular nets, $f > 1$. One can realize such nets on the surfaces of curvature -1 . Let us consider f regular k -gons on the hypobolic plane with angles equal $2\pi/3$ and label the sides according these words. Such partitions of the hyperbolic plane with an action of the fundamental group of M_g^2 according the following words gives us examples of surfaces with an "exotic" local minimal net.

Let f be 2 and k be 12. Then there is the following example

$$e^{-1}abca^{-1}db^{-1}c^{-1}d^{-1}elm^{-1},$$

$$mnfghf^{-1}ig^{-1}h^{-1}i^{-1}n^{-1}l^{-1}.$$

Let $f = 3$ and $k = 10$:

$$a_1a_4a_5a_7a_1^{-1}a_8a_5^{-1}a_{14}^{-1}a_{10}^{-1}a_9^{-1}$$

$$a_{14}a_1^{-1}a_7^{-1}a_8^{-1}a_9a_{13}^{-1}a_{15}^{-1}a_{18}^{-1}a_{19}^{-1}a_{11}^{-1}$$

$$a_{10}a_{11}a_{12}a_{15}a_{16}a_{18}a_{12}^{-1}a_{19}a_{16}^{-1}a_{13}$$

Let $f = 4$ and $k = 9$:

$$a_1a_4a_5a_2a_3a_{15}^{-1}a_{18}^{-1}a_{19}^{-1}a_9^{-1}$$

$$a_2^{-1}a_6a_7a_1^{-1}a_8a_6^{-1}a_5^{-1}a_{11}^{-1}a_{10}$$

$$a_{11}a_4^{-1}a_7^{-1}a_8^{-1}a_9a_{12}a_{15}a_{16}a_{14}^{-1}$$

$$a_{14}a_{17}a_{18}a_{12}^{-1}a_{19}a_{17}^{-1}a_{16}^{-1}a_3^{-1}a_{10}$$

Let $f = 6$ and $k = 8$:

$$a_1a_4a_5a_2a_3a_{20}a_{10}^{-1}a_9^{-1}$$

$$a_2^{-1}a_6a_7a_1^{-1}a_8a_6^{-1}a_5^{-1}a_{13}^{-1}$$

$$a_{13}a_4^{-1}a_7^{-1}a_8^{-1}a_9a_{14}a_{21}a_3^{-1}$$

$$a_{10}a_{11}a_{12}a_{15}a_{16}a_{23}^{-1}a_{22}^{-1}a_{14}^{-1}$$

$$a_{17}a_{18}a_{12}^{-1}a_{19}a_{17}^{-1}a_{16}^{-1}a_{24}a_{23}$$

$$a_{15}^{-1}a_{18}^{-1}a_{19}^{-1}a_{11}^{-1}a_{20}a_{21}a_{22}a_{24}^{-1}$$

Let $f = 12$ and $k = 7$:

$$a_1a_2a_{20}a_{21}a_{22}a_{10}^{-1}a_9^{-1}$$

$$a_3a_4a_5a_{23}a_{24}a_{25}a_{20}^{-1}$$

$$\begin{aligned}
& a_6 a_7 a_3^{-1} a_2^{-1} a_{26} a_{27} a_{23}^{-1} \\
& a_1^{-1} a_8 a_6^{-1} a_5^{-1} a_{29} a_{28}^{-1} a_{26}^{-1} \\
& a_4^{-1} a_7^{-1} a_8^{-1} a_9 a_{30} a_{31}^{-1} a_{29}^{-1} \\
& a_{10} a_{11} a_{12} a_{13} a_{33}^{-1} a_{32}^{-1} a_{30}^{-1} \\
& a_{14} a_{15} a_{16} a_{34} a_{35} a_{36} a_{33} \\
& a_{17} a_{18} a_{14}^{-1} a_{13}^{-1} a_{37} a_{38} a_{34}^{-1} \\
& a_{12}^{-1} a_{19} a_{17}^{-1} a_{16}^{-1} a_{40} a_{39} a_{37}^{-1} \\
& a_{15}^{-1} a_{18}^{-1} a_{11}^{-1} a_{22}^{-1} a_{41} a_{40}^{-1} \\
& a_{21}^{-1} a_{25}^{-1} a_{42} a_{35}^{-1} a_{38}^{-1} a_{39}^{-1} a_{41}^{-1} \\
& a_{42}^{-1} a_{24}^{-1} a_{27}^{-1} a_{28} a_{31}^{-1} a_{32} a_{36}^{-1}
\end{aligned}$$

References

- [1] F.J. Almgren, Jr. and J.E. Taylor, The Geometry of soap films and soap bubbles, Scientific American, 1976
- [2] A. Heppes, Isogonal sphärischen netze, Ann. Univ. Sci. Budapest Eötvös Sect. Math. **7** (1964), 41-48.
- [3] A.T. Fomenko, The Plateau Problem (New York, Gordon and Breach, 1989)
- [4] A.T. Fomenko, Variational Problems in Topology (New York, Gordon and Breach, 1990)
- [5] A.Vdovina Constructing of orientable Wicks forms and estimation of their number. Commun. in Algebra, V.23 (1995), 9, 3205-3222
- [6] Vdovina A.A., Selivanova E.N. Local minimal Networks on surfaces of constant negative curvature, Vestnik MGU, (1997), **6**, 15-17 (Russian)
- [7] T.R. Walsh, A.B. Lehman Counting Rooted maps by Genus, J. Comb. Theory, (B), 13, 192-218 (1972)